

UNCLASSIFIED

AD 410110

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

N-63-4-3

CATALOGED BY DDC  
AS AD No. 410110

OFFICE OF NAVAL RESEARCH

Contract Nonr 3395(00)

Task No. NR 356-431

TECHNICAL REPORT NO. 18

Chemistry of Decaborane-Phosphorus Compounds. IV.  
Monomeric, Oligomeric and Cyclic Phosphino-Carboranes.

by

Roy P. Alexander and Hansjuergen Schroeder

Prepared for Publication

in

Inorganic Chemistry

DDC  
REFILED  
JUN 17 1963

Olin Mathieson Chemical Corporation

Organics Division

New Haven, Connecticut

JUNE 1963

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

410110

CONTRIBUTION FROM THE ORGANICS DIVISION

OLIN MATHIESON CHEMICAL CORPORATION

Chemistry of Decaborane-Phosphorus Compounds. IV.  
Monomeric, Oligomeric and Cyclic Phosphino-Carboranes.<sup>1,2</sup>


By Roy P. Alexander and Hansjuergen Schroeder

Received.....

Dilithiocarborane,  $B_{10}H_{10}C_2Li_2$ , was allowed to react with chlorophosphines containing one, two, and three chlorine atoms.  $(C_6H_5)_2PCl$  and  $C_6H_5PCl_2$  gave the expected disubstituted compounds,  $B_{10}H_{10}C_2[P(C_6H_5)_2]_2$  and  $B_{10}H_{10}C_2(C_6H_5P\cdot Cl)_2$ , respectively.  $B_{10}H_{10}C_2Li_2$  and  $PCl_3$ , however, formed a compound having two phosphorus atoms and two carborane nuclei in a six membered ring. This unique ring system also arose from the reaction of  $B_{10}H_{10}C_2Li_2$  with  $B_{10}H_{10}C_2-(C_6H_5P\cdot Cl)_2$ . Subsequent reactions of such molecules with ammonia and  $NaN_3$  are described. The cyclic bis-azide,  $(B_{10}H_{10}C_2P\cdot N_3)_2$ , and  $p-[P(C_6H_5)_2]_2C_6H_4$  reacted to form a PNP bonded oligomer.

In previous papers of this series we reported the synthesis of bifunctional decaborane-phosphine monomers<sup>3</sup> and their conversion into linear polymers<sup>1</sup>. The high-thermal and hydrolytical stability of PNP bonded decaborane-phosphine polymers<sup>1</sup>, especially of compound I, made them an obvious object for further investigation.

- 
- (1) Preceding communication: H. Schroeder, J. R. Reiner, and T. A. Knowles, *Inorg. Chem.*, 2, 393 (1963).
  - (2) Presented before the Inorganic Division of The American Chemical Society at the 145th Meeting at New York, N.Y., Sept. , 1963.
  - (3) H. Schroeder, J. R. Reiner, and T. L. Heying, *Inorgan. Chem.*, 1, 618 (1962).




Since it was established<sup>1</sup> that compound I lost hydrogen from the decaborane nuclei under crosslinking at 340°, a logical extension of this work with regard to improving heat stability was the replacement of those hydrogens by atoms more firmly bonded to boron such as chlorine and fluorine. Initial attempts to chlorinate the starting monomer  $\text{B}_{10}\text{H}_{10}[(\text{C}_6\text{H}_5)_2\text{P}\cdot\text{Cl}]_2$  met with failure. However, before a more thorough halogenation study of compounds containing the  $\text{B}_{10}\text{H}_{12}$  unit was conducted, attention was turned to the easily chlorinated carborane<sup>4</sup> moiety instead.

Interest in this laboratory has been focused recently on the chlorination of carborane resulting in polychlorocarboranes such as  $\text{B}_{10}\text{Cl}_{10}\text{C}_2\text{H}_2$ .<sup>5</sup> The potential utility of the chlorinated carborane nucleus for the formation of PNP bonded carborane polymers led us to initiate an investigation aimed at the preparation of difunctional phosphorus containing carborane and polychlorocarborane derivatives. The present study describes the first part of our research dealing with the synthesis of chlorophosphino-carboranes followed by the application of the PNP polymer scheme to those compounds obtained containing the unchlorinated carborane nucleus.

The ability of carborane to undergo direct lithiation upon treatment with n-butyllithium<sup>6</sup> provided dilithiocarborane, an appropriate starting compound for reactions with various chloro-


- 
- (4) For brevity, the term carborane is used for 11,12-dicarba-dodecaborane; for structure of carborane see reference 4.
  - (5) H. Schroeder, T. L. Heying, and J. R. Reiner, *Inorg. Chem.*, ..... (1963).
  - (6) T. L. Heying, J. W. Ager, Jr., S. L. Clark, R. P. Alexander, S. Papetti, J. A. Reid, and S. I. Trotz, submitted for publication.

phosphines. In a prototype experiment, methathesis was found to occur between dilithiocarborane and diphenylchlorophosphine to give bis(diphenylphosphino)-carborane.<sup>7</sup>



Having demonstrated the fundamental reaction, the dilithio salt was reacted with phenyldichlorophosphine. As expected, two molar equivalents of  $\text{C}_6\text{H}_5\text{PCl}_2$  reacted with one equivalent of  $\text{B}_{10}\text{H}_{10}\text{C}_2\text{Li}_2$  to form the desired bis-chloro compound  $\text{B}_{10}\text{H}_{10}\text{C}_2(\text{C}_6\text{H}_5\text{P}\cdot\text{Cl})_2$ .

To determine the reactivity of the phosphorus bound chlorine atoms, ammonia was passed, at room temperature, into a solution of  $\text{B}_{10}\text{H}_{10}\text{C}_2(\text{C}_6\text{H}_5\text{P}\cdot\text{Cl})_2$  in benzene. Ammonolysis occurred, but instead of the expected bis-amino compound  $\text{B}_{10}\text{H}_{10}\text{C}_2[\text{P}(\text{C}_6\text{H}_5)\text{NH}_2]_2$ , a cyclic diphospha(III)azane,  $\text{B}_{10}\text{H}_{10}\text{C}_2[(\text{PC}_6\text{H}_5)_2\text{NH}]$ , was recovered in 70% yield. The infrared spectrum was consistent with this formulation in showing only a single N-H stretching absorption at  $3\mu$ .




On the other hand, treatment of  $\text{B}_{10}\text{H}_{10}\text{C}_2(\text{C}_6\text{H}_5\text{P}\cdot\text{Cl})_2$  with sodium azide in ethanol at  $0^\circ$  gave the desired bis-azide,  $\text{B}_{10}\text{H}_{10}\text{C}_2(\text{C}_6\text{H}_5\text{P}\cdot\text{N}_3)_2$ , m.p.  $126-128^\circ$  dec. In benzene solution the compound undergoes decomposition at approximately  $40^\circ$  with evolution of nitrogen yielding a product, the analysis of which is in general agreement with the formula  $(=\text{NP}(\text{C}_6\text{H}_5)-\text{CB}_{10}\text{H}_{10}\text{C}-(\text{C}_6\text{H}_5)\text{PN}=\text{})_x$ . The decomposition is promoted by the addition of

(7) The notation  $\underset{\text{B}_{10}\text{H}_{10}}{\text{C}}$  will be employed throughout this publication to designate the structure of the carboranyl grouping.


a phosphine such as  $(C_6H_5)_3P$ , since in their presence nitrogen evolution begins at  $10^\circ$  to  $20^\circ$ . The phosphines do not interact with this phosphorus(III)bis-azide, a finding in sharp contrast to the easy formation of phosphorus-nitrogen double bonds<sup>1</sup> observed in the reaction of  $B_{10}H_{10}[(C_6H_5)_2P \cdot N_3]_2$ , a compound having pentavalent phosphorus, with phosphines.

In view of these results it appeared of interest to investigate next the reaction of  $B_{10}H_{10}C_2Li_2$  with  $PCl_3$ . It was found that equimolar amounts of these reactants form a product which analyzes as  $(B_{10}H_{10}C_2PCl)_x$ . On the evidence of an accurate cryoscopic molecular weight determination corroborated by mass spectral analysis, a cyclic structure containing two carborane moieties and two phosphorus atoms in the ring was assigned for this reaction product. Another example of this diphosphacyclohexane system was obtained by reaction of  $B_{10}H_{10}C_2(C_6H_5 \cdot P \cdot Cl)_2$  with  $B_{10}H_{10}C_2Li_2$  to give the diphenyl substituted derivative. However, the dichloro-compound has resisted attempts to convert to the diphenyl compound by either Friedel-Crafts, Grignard, or aryl-lithium procedures.



In a comparison of the carborane infrared spectrum with those of  $B_{10}H_{10}C_2(C_6H_5P \cdot Cl)_2$  and  $(B_{10}H_{10}C_2P \cdot Cl)_2$  (see Fig. 1) it is of interest to note that not only does the C-H stretching frequency disappear, but also the carborane and chlorocarborane characteristic absorptions<sup>5</sup> between 8 and 9  $\mu$  are not present. The simplicity of the spectrum of the cyclic bischloro-compound also indicates high symmetry for this molecule.

Treatment of the cyclic bis-chloro compound with ammonia and with sodium azide, respectively, afforded in good yields the cyclic bis-amino compound,  $(B_{10}H_{10}C_2P \cdot NH_2)_2$ , and the cyclic bis-azide,  $(B_{10}H_{10}C_2P \cdot N_3)_2$ . Unlike the preceding bis-azide,  $B_{10}H_{10}C_2(C_6H_5P \cdot N_3)_2$ , it proved to be stable in refluxing benzene and reacted with triphenylphosphine according to the following equation:




This is the first example of a phosphorus(III)azide reacting with a phosphine to form a phosphineimino compound.


Upon extension of this reaction to the use of a diphosphine following the PNP-polymer pattern, equimolar amounts of  $(B_{10}H_{10}C_2P \cdot N_3)_2$  and  $p-[(C_6H_5)_2P]_2C_6H_4$  reacted with evolution of nitrogen to form compound II.



FIGURE 1.

INFRARED SPECTRA OF CARBORANE, BIS(CHLOROPHENYLPHOSPHINO)-CARBORANE,  
AND DIMERIC CHLOROPHOSPHA (III) CARBORANE (Kel-F and Nujol Mulls)





This reaction was repeated several times and in each instance the elemental analysis was precise for the structure shown, and molecular weight determination gave  $x$  a value 3 ( $\pm 5\%$ ). A rigorous structural determination of this apparently trimeric substance is in progress.

EXPERIMENTAL<sup>8</sup>

Starting Materials.-  $B_{10}H_{10}C_2H_2$  was prepared as described in ref. 6 in n-propyl ether (Eastman Kodak) from resublimed  $B_{10}H_{14}$ , diethylsulfide (Matheson, Coleman & Bell) and acetylene which was purified by standard procedures.<sup>9</sup> Diphenylchlorophosphine was obtained from Victor Chemical Company.

Molecular Weights.- These were obtained in benzene either using a Mechrolab Osmometer, Model 301, or by cryoscopic determination.

Dilithiocarborane<sup>6</sup>.- A 100 ml. three-neck flask fitted with a magnetic stirrer, a nitrogen inlet and outlet, and an addition funnel, is purged with nitrogen and then charged with a solution of butyllithium (0.066 mole) in ethyl ether (approx. 60 ml.). With ice-cooling a solution of carborane (4.33 g., 0.3 mole) in ethyl ether (20 ml.) is added over a period of 10 minutes, and the resulting white slurry is stirred under nitrogen and ice-cooling for another 30 minutes. Ether and excess butyllithium are then filtered from the precipitate through a coarse sintered-glass filter stick. The remaining  $B_{10}H_{10}C_2Li_2$  is washed with ethyl ether (50 ml.) in two portions, dried under vacuum and then stored in an inert atmosphere.

Bis(diphenylphosphino)carborane.- A solution of diphenylchlorophosphine (14.6 g., 0.066 mole) in ether (15 ml.) was added, with ice-cooling, within 25 minutes to a dilithiocarborane slurry in ether prepared from 4.33 g. (0.3 mole) of carborane as described

- 
- (8) Melting points are uncorrected; elemental analyses were by R. Culmo, A. C. Mayer, and J. D. Giunta of the Olin Mathieson Microchemical Section; mass spectral analyses were obtained and interpreted by Herman Hoberecht.
- (9) We thank Mr. R. W. Blundon for preparing a continuing supply of carborane for this research.

in the preceding example. The slightly yellow mixture was stirred for 20 minutes at 20°, then refluxed for 30 minutes and, after cooling to 10°, hydrolyzed with 20 ml. of water. The insoluble reaction product was recovered by filtration, dried in vacuo and recrystallized from petroleum ether (b.p. 60-110°) to give 9.0 g. (58.5%) of  $B_{10}H_{10}C_2[P(C_6H_5)_2]_2$ , m.p. 219°.

Anal. Calcd. for  $C_{26}H_{30}B_{10}P_2$  (512.7): C, 60.91; H, 5.90; B, 21.11; P, 12.09. Found: C, 61.15; H, 6.23; B, 20.63; P, 11.68.

Bis(chlorophenylphosphino)carborane.— A dilithiocarborane (0.173 mole) slurry in ether prepared from 25 g. of carborane was slowly added to a cooled solution of phenyldichlorophosphine (77.5 g., 0.43 mole) in ether (600 ml.). The mixture was stirred at room temperature for 90 minutes, then refluxed for 1 hour. Upon evaporation of the ether a yellow solid remained which was extracted with hot petroleum ether (800 ml., b.p. 60-110°). After concentration and cooling, 43.7 g. (58%) of  $B_{10}H_{10}C_2(C_6H_5P\cdot Cl)_2$ , m.p. 172-174°, precipitated from the solution.

Anal. Calcd. for  $C_{14}H_{20}B_{10}Cl_2P_2$  (429.4): C, 39.16; H, 4.70; B, 25.20; Cl, 16.51; P, 14.43. Found: C, 38.87; H, 4.67; B, 24.95; Cl, 16.81; P, 14.18; mol. weight: 420.1.

C,C'-Bis phenylphospha(III) -azanyl(1,3)-carborane.— Gaseous ammonia was bubbled into a solution of  $B_{10}H_{10}C_2(C_6H_5P\cdot Cl)_2$  (4.3 g., 0.01 mole) in anhydrous benzene (100 ml.) for 30 minutes. The temperature was maintained at 20° by external cooling. The reaction mixture was then evaporated to dryness, the solid residue was refluxed with 150 ml. of hot petroleum ether (b.p. 60-110°). After filtration, concentration, and cooling, the imino-compound precipitated and was recrystallized

from petroleum ether. Yield of  $B_{10}H_{10}C_2[(PC_6H_5)_2NH]$ : 2.6 g. (70%); m.p. 222-24°.

Anal. Calcd. for  $C_{14}H_{21}B_{10}NP_2$  (373.5): C, 45.02; H, 5.67; B, 28.97; N, 3.75; P, 16.59. Found: C, 45.00; H, 5.82; B, 28.46; N, 3.72; P, 16.65.

Bis(azidophenylphosphino)carborane.- A mixture of  $B_{10}H_{10}C_2(C_6H_5P \cdot Cl)_2$  (10.7 g., 0.025 mole), sodium azide (6.5 g., 0.1 mole), and ethanol (150 ml.), was stirred with ice-cooling for 1 hour. The solid reaction products were recovered by filtration and thoroughly washed with water. The remaining bis-azide was dried in vacuo over diphosphorus pentoxide; m.p. 126-28° (dec.); yield: 9.1 g. (82%).

Anal. Calcd. for  $C_{14}H_{20}B_{10}N_6P_2$  (442.5): C, 38.00; H, 4.56; B, 24.45; N, 18.99; P, 14.00. Found: C, 38.29; H, 4.60; B, 24.55; N, 17.80; P, 13.85.

Pyrolysis: A solution of 6 g. of  $B_{10}H_{10}C_2(C_6H_5P \cdot N_3)_2$  in 125 ml. dry benzene was slowly warmed. At a bath temperature of approximately 50° gas evolution started. After keeping the mixture for 90 minutes at 55°, it was cooled to 20° and the precipitate formed (2.5 g.) was recovered by filtration. It did not show the characteristic azide absorption at  $4.8\mu$ ; m.p. 170-180°.

Anal. Calcd. for  $(C_{14}H_{20}B_{10}N_2P_2)_x$  (386.4): C, 43.51; H, 5.22; B, 28.00; N, 7.23; P, 16.03. Found: C, 41.93; H, 4.82; B, 26.44; N, 6.22; P, 14.89.

Dimeric Chlorophospha(III)carborane.- A dilithiocarborane (0.139 mole) slurry in ether (500 ml.) prepared from 20 g. of carborane was added in small portions, with ice-cooling, to a stirred solution of phosphorus trichloride (41.1 g., 0.3 mole) in ether (1000 ml.). The mixture immediately became purple and a colorless precipitate began to form. After stirring for 20 hours at 20°, the mixture had turned blue and was evaporated to dryness. The green solid residue was extracted with 300 ml. of

Dimeric Triphenylphosphineiminophospha(III)carborane.- To a mixture of  $(B_{10}H_{10}C_2P.N_3)_2$  (2.15 g., 0.005 mole) and triphenylphosphine (2.62 g., 0.01 mole) was added, in one portion, cold benzene (50 ml.). Solution was effected accompanied by vigorous foaming. After 25 minutes, gas evolution subsided and the solution was gradually heated to reflux temperature over a 30 minute-period. The hot reaction mixture was filtered and cooled to  $10^\circ$ . The colorless reaction product precipitated and was washed with petroleum ether. Yield of  $[(C_6H_5)_3P=NPC_2B_{10}H_{10}]_2$ : 2.4 g. (53.5%), m.p.  $311-12^\circ$ .

Anal. Calcd. for  $C_{40}H_{50}B_{20}N_2P_4$  (899.2): C, 53.43; H, 5.61; B, 24.07; N, 3.12; P, 13.78. Found: C, 53.89; H, 5.92; B, 24.08; N, 3.54; P, 13.73.

Reaction of Dimeric Azidophospha(III)carborane with 1,4-Bis-(diphenylphosphino)-benzene.- Addition of cold benzene (75 ml.) to a mixture of  $(B_{10}H_{10}C_2P.N_3)_2$  (3.45 g., 0.008 mole) and p- $[(C_6H_5)_2P]_2C_6H_4$  (3.55 g., 0.008 mole) produced, with evolution of nitrogen, a clear, purple solution. Stirring at  $20^\circ$  for 1 hour followed by refluxing for 2 hours changed the color to pale-blue. After filtration from a small amount of solid, the filtrate was poured into 500 ml. of petroleum ether. A solid separated, was recovered by filtration and purified by washing with hot petroleum ether. Yield of product: 4.6 g. (70%); m.p.  $235-50^\circ$ .

Anal. Calcd. for  $C_{34}H_{44}B_{20}N_2P_4$  (821.0): C, 49.74; H, 5.40; B, 26.36; N, 3.41; P, 15.09. Found: C, 50.20; H, 5.65; B, 26.06; N, 3.05; P, 14.59; molecular weight: 2367(experiment 1), 2585(experiment 2), 2498(experiment 3).

Dimeric Phenylphospha(III)carborane.- A solution of  $B_{10}H_{10}C_2(C_6H_5P.Cl)_2$  (6.9 g., 0.016 mole) in ether (200 ml.) was added, with stirring, to an ice-cooled dilithiocarborane (0.016) slurry in ether (100 ml.) prepared from 2.32 g. of carborane. After stirring the mixture for 12 hours at room temperature, it was filtered to give 3.6 g. of insoluble residue. Extraction with hot benzene afforded 2.0 g. of product which was recrystallized from benzene to give 1.6 g. (20%) of  $(B_{10}H_{10}C_2P.C_6H_5)_2$ , m.p. 356-358°.

Anal. Calcd. for  $C_{16}H_{30}B_{20}P_2$  (500.8): C, 38.38; H, 6.04; B, 43.21; P, 12.37. Found: C, 38.28; H, 5.79; B, 42.36; P, 11.93.

ACKNOWLEDGMENT.- We are grateful to Dr. T. L. Heying for stimulating discussions. This work was supported by the Office of Naval Research.

# TECHNICAL REPORT DISTRIBUTION LIST

CONTRACTOR OLINGATHURSON CHEMICALS CORPORATION

NR NO. 356-431

CONTRACT NUMBER Hour 3355(00)

DATE 15 FEB 1962

	<u>NO. COPIES</u>		<u>NO. COPIES</u>
Commanding Officer Office of Naval Research Branch Office The John Crerar Library Building 100 East Randolph Street Chicago 1, Illinois	(1)	Air Force Office of Scientific Research (SRC-E) Washington 25, D. C.	(1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York	(1)	Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D. C. Attn: Technical Information Office Branch 012	(1)
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena 1, California	(1)	Office, Chief of Research & Development Department of the Army Washington 25, D. C. Attn: Physical Sciences Division	(1)
Commanding Officer Office of Naval Research Branch Office Box 39 Navy #100 Fleet Post Office New York, New York	(7)	Chief, Bureau of Ships Department of the Navy Washington 25, D. C. Attn: Code 342A Code 634C	(2) (1)
Director, Naval Research Laboratory Washington 25, D. C. Attn: Technical Information Officer Chemistry Division	(6) (2)	Chief, Bureau of Naval Weapons Department of the Navy Washington 25, D. C. Attn: Technical Library Code RRM-3	(3) (1)
Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 425	(2)	ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia	(10)
DDR&E Technical Library Room 3C-128, The Pentagon Washington 25, D. C.	(1)	Director of Research U.S. Army Signal Research & Development Laboratory Fort Monmouth, New Jersey	(1)
Technical Director Research & Engineering Division Office of the Quartermaster General Department of the Army Washington 25, D. C.	(1)	Naval Radiological Defense Laboratory San Francisco 24, California Attn: Technical Library	(1)
Research Director Clothing & Organic Materials Division Quartermaster Research & Engineering Command U. S. Army Natick, Massachusetts	(1)	Naval Ordnance Test Station China Lake, California Attn: Head, Chemistry Division Code 40 Code 50	(1) (1)

REVISED 1 FEB 1962



TECHNICAL REPORT DISTRIBUTION LIST

Page 2

CONTRACTOR OLYN MATHIESON CHEMICALS CORPORATION

NR NO. 356-431

CONTRACT NUMBER Monr 3395(00)

DATE 15 FEB. 1962

<u>NO. COPIES</u>	<u>NO. COPIES</u>
Commanding Officer Army Research Office Box CM, Duke Station Durham, North Carolina Attn: Scientific Synthesis Office (1)	Dr. Albert Lightbody Naval Ordnance Laboratory White Oak, Maryland (1)
Brookhaven National Laboratory Chemistry Department Upton, New York (1)	Commanding Officer and Director U. S. Naval Civil Engineering Lab. Port Hueneme, California Attn: Chemistry Division (1)
Atomic Energy Commission Division of Research Chemistry Programs Washington 25, D. C. (1)	Dr. A. L. Powell Office of Naval Research Branch Office 495 Summer Street Boston 10, Mass. (1)
Atomic Energy Commission Division of Technical Information Extension Post Office Box 62 Oak Ridge, Tennessee (1)	Aeronautical Systems Division ASRCNP Wright-Patterson Air Force Base Ohio (1)
U.S. Army Chemical Research and Development Laboratories Technical Library Army Chemical Center, Maryland (1)	Dr. H. C. Clark Department of Chemistry University of British Columbia Vancouver, British Columbia, Canada (1)
Office of Technical Services Department of Commerce Washington 25, D. C. (1)	Dr. E. G. Rochow Department of Chemistry Harvard University Cambridge 38, Mass. (1)
Dr. P. A. Miller Office of Naval Res. Br. Off. 1000 Geary Street San Francisco 9, Calif. (1)	Dr. John E. Leffler Department of Chemistry Florida State University Tallahassee, Fla. (1)
Dr. C. Haber Naval Ordnance Laboratory Corona, California (1)	Dr. William N. Lipscomb Department of Chemistry Harvard University Cambridge, Mass. (1)
Dr. Porter W. Erickson Chemistry Research Department Non-Metallic Materials Division Naval Ordnance Laboratory White Oak, Maryland (1)	Dr. T. D. Parsons Department of Chemistry Oregon State College Corvallis, Oregon (1)

TECHNICAL REPORT DISTRIBUTION LIST

Page 3

CONTRACTOR OLIN MATHIESON CHEMICALS CORPORATION

HR NO. 356-431

CONTRACT NUMBER Nonr 3395(00)

DATE 15 FEB 1962

	<u>NO. COPIES</u>		<u>NO. COPIES</u>
Dr. L. F. Rahn Princeton University Princeton Plastics Laboratory Princeton, New Jersey	(1)	Aircraft Industries Association 7660 Beverly Boulevard Los Angeles 36, Calif. Attn: Mr. H. D. Moran	(10)
Dr. A. V. Tobolsky Department of Chemistry Princeton University Princeton, New Jersey	(1)	American Potash & Chem. Corp. 201 W. Washington Blvd. Whittier, California Attn: Dr. W. S. Emerson	(2)
Dr. R. S. Stein Department of Chemistry University of Massachusetts Amherst, Massachusetts	(1)	U. S. Borax Research Corp. Attn: Dr. Carl Randolph Anaheim, California	(2)
Dr. S. Young Tyree, Jr. Department of Chemistry University of North Carolina Chapel Hill, North Carolina	(1)	General Electric Company Research Laboratory P. O. Box 1088 Schenectady, New York Attn: Dr. J. R. Elliot	(2)
Dr. J. C. Bailar, Jr. Department of Chemistry University of Illinois Urbana, Illinois	(2)	Dr. P. D. George General Electric Company General Engineering Lab. Schenectady, New York	(2)
Dr. L. F. Audrieth Department of Chemistry University of Illinois Urbana, Illinois	(2)	Dr. Hans B. Jonassen Department of Chemistry Tulane University New Orleans 15, Louisiana	(1)
Dr. A. B. Burg Department of Chemistry University of Southern California Los Angeles 7, Calif.	(2)	Dr. Henry Taube Department of Chemistry Stanford University Stanford, Calif.	(1)
Dr. Riley Schaeffer Department of Chemistry Indiana University Bloomington, Indiana	(1)	Plastics Technical Evaluation Center Picatinny Arsenal Dover, N. J.	(1)
Dr. T. G. Fox, Jr., Director of Research Mellon Institute 4400 Fifth Avenue Pittsburgh 13, Pennsylvania	(1)	Dr. G. Barth-Wehrenalp, Director Inorganic Research Department Pennsalt Chemicals Corporation Box 4388 Philadelphia 18, Penna.	(2)

TECHNICAL REPORT DISTRIBUTION LIST

Page 4

CONTRACTOR OLIN MATHIESON CHEMICALS CORPORATION

NR NO. 356-431

CONTRACT NUMBER Nonr 3395(00)

DATE 1 FEB 1962

	<u>NO. COPIES</u>		<u>NO. COPIES</u>
Dr. M. M. Chamberlain Department of Chemistry Western Reserve University Cleveland, Ohio	(1)	Dr. D. A. Brown Department of Chemistry University College Upper Merrion Street Dublin, Ireland	(1)
Dr. D. C. Bradley Department of Chemistry University of Western Ontario London, Canada	(1)	Dr. Albert Cotton Department of Chemistry Massachusetts Institute of Technology Cambridge 39, Mass.	(1)
Dr. T. P. Onak Department of Chemistry Los Angeles State College Los Angeles, California	(1)	Dr. Robert R. Holmes Department of Chemistry Carnegie Institute of Technology Pittsburgh 13, Penn.	(1)
Dr. M. J. S. Dewar Department of Chemistry University of Chicago Chicago, Illinois	(1)	Prof. L. A. Bigelow Department of Chemistry Duke University Durham, N. C.	(1)
Dr. M. S. Cohen Thiokol Chemical Corporation Reaction Motors Division Denville, New Jersey	(1)	Dr. William T. Miller Department of Chemistry Cornell University Ithaca, New York	(1)
Inspector of Naval Material 181 Middle Street Bridgeport 3, Conn.	(1)	Prof. Jack Hine Department of Chemistry Georgia Institute of Technology Atlanta 3, Georgia	
Dr. George F. Huff, Vice President Research and Development Callery Chemical Company Callery, Penn.	(1)	Prof. Herbert C. Brown Department of Chemistry Purdue University Lafayette, Indiana	(1)
Dr. Joyce J. Kaufman RIAS 7212 Bellona Avenue Baltimore 12, Maryland	(1)	Dr. Phillip S. Skell Department of Chemistry Pennsylvania State University University Park, Penn.	(1)
Dr. Arthur Finch Royal Holloway College University of London London, England	(1)	Dr. B. D. Post Polytechnic Institute of Brooklyn Brooklyn 1, New York	(1)

TECHNICAL REPORT DISTRIBUTION LIST

Page 5

CONTRACTOR OLIN MATHIESON CHEMICALS CORPORATION

NR NO. 356-431

CONTRACT NUMBER Nonr 3395(00)

DATE 15 FEB, 1962

NO. COPIES

Dr. R. T. Holzmann  
Advanced Research Projects Agency  
Washington 25, D. C. (1)

Commander  
Ordnance Corps  
Rock Island Arsenal  
Rock Island, Illinois  
Attn: ORDBC 9320-ZTO (1)

New York Naval Shipyard  
Material Laboratory  
Brooklyn 1, New York  
Attn: Mr. B. B. Simms (1)

Bureau of Ships,  
Department of the Navy  
Washington 25, D. C.  
Code 660L  
Attn: Mr. E. J. Hrycklewicz (1)

Dr. M. F. Hawthorne  
Department of Chemistry  
University of California, Riverside  
Riverside, California (1)

Dr. Roald Hoffman  
Department of Chemistry  
Harvard University  
Cambridge, Massachusetts (1)

Mr. J. A. Kies  
Code 6210  
Naval Research Laboratory  
Washington 25, D. C. (1)

Mr. E. J. Kohn  
Code 6110  
Naval Research Laboratory  
Washington 25, D. C. (1)

Dr. R. B. Fox (1)  
Mr. J. E. Cowling (1)  
Dr. A. L. Alexander (1)  
Dr. D. L. Venezky (1)  
Code 6120  
Naval Research Laboratory  
Washington 25, D. C.

Dr. O. Williams  
National Science Foundation  
Washington 25, D. C. (1)

Monsanto Research Corporation  
1515 Nicholas Road  
Dayton, Ohio  
Attn: Librarian (1)

Monsanto Research Corporation  
Everett Station  
Boston 49, Massachusetts  
Attn: Librarian

The Dow Chemical Company  
ARPA Laboratory  
1710 Building  
Midland, Michigan (1)

Naval Ordnance Test Station  
China Lake, California  
Attn: Code 4544 (Dr. Kaufman) (1)  
Code 5557 (Mr. S. H. Herzog)  
(Mr. R.J. Landry) (1)